






NATIONAL
WEATHER
SERVICE

UFS-S2S Prototype 5 Benchmark Evaluation Summary





Technical and Science Contributors



- **NCEP/EMC:** Jessica Meixner, Lydia Stefanova, Jiande Wang, Denise Worthen, Shrinivas Moorthi, Bin Li, Jun Wang, Jian Kuang, Partha Bhattacharjee, Robert Grumbine, Wei Li, Weizhong Zheng, Suranjana Saha, Huug van den Dool, Xingren Wu, Roberto Padilla, Fanglin Yang, Arun Chawla, Avichal Mehra, Vijay Tallapragada
- **ESRL:** Shan Sun, Rainer Bleck, Benjamin W. Green
- **NESII:** Robert Oehmke, Gerhard Theurich
- **CICE Consortium:** Tony Craig
- **GFDL:** Rusty Benson, Brandon Reichl, Stephen M. Griffies, Robert Hallberg, Alistair Adcroft
- **UCAR/NCAR:** Mariana Vertenstein, Rocky Dunlap, Dave Bailey, Alper Altuntas, Gustavo Marques, Ufuk Turuncoglu, Dan Rosen

For additional information, please contact Lydia.B.Stefanova@noaa.gov



Prototype Overview

	Initial Conditions					
	FV3 GFS	MOM6	CICE	WW3	Ice Model	Mediator
UFS_p1	CFSR	CFSR	CFSR	n/a	CICE5	NEMS
UFS_p2	CFSR	CPC 3Dvar	CFSR	n/a	CICE5	NEMS
UFS_p3.1	CFSR	CPC 3Dvar	CPC ice analysis	n/a	CICE5	NEMS
UFS_p4	CFSR	CPC 3Dvar	CPC ice analysis	Generated with CFS forcings	CICE5	NEMS
UFS_p5	CFSR	CPC 3DVar	CPC ice analysis	Generated with CFS forcings	CICE6	CMEPS

- Physics settings are kept the same as much as possible except for bug fixes.
- **Physics tuning is reserved for after engineering is completed**
- The aim of prototype evaluations to date is to **establish baseline skill and ensure technical consistency between prototypes**



Anomaly Correlation Skill for Select Variables/Domains

Weeks 1, 2, 3&4





Notes on the methodology used here



Results are shown for both raw and systematic-error-corrected (sec) anomalies.



The systematic error correction is applied by fitting and removing a smoothly interpolated climatology to both the forecast and observation data. This climatology is produced by fitting the 7 year time series (168 elements, 1st and 15th of each month) to a sine wave of period 365.24 days, plus three harmonics.



This is done for each gridpoint and variable separately for both for forecasts (as a function of lead) and verifying data.



All forecasts were bias corrected in exactly the same manner.





Summary of findings



- Skill improvement from SEC depends on field, domain, lead
- Prototype 5 AC skill is on par with preceding prototypes and above that of the operational CFSv2, with the exception of weeks 3&4 CONUS T2m. The latter is tentatively attributed to absence of lake ice in Prototype 5 which will be corrected in Prototype 6.
- Annual SEC-corrected AC skill scores for P5:
 - Week 1
 - 0.9+ for Z500, Nino3.4 SST
 - 0.85+ for Tropical SST and CONUS T2m
 - 0.55+ for Nino3.4 Precipitation and CONUS Precipitation
 - Week 2
 - ~0.9 for Nino3.4 SST, ~0.8 for Tropical SST
 - ~0.5-0.6 for Z500, 0.4-0.5 for Nino3.4 Precipitation, CONUS T2m
 - ~0.2 for CONUS Precipitation
 - Weeks 3&4
 - ~0.9 for Nino3.4 SST, 0.7-0.75 for Tropical SST
 - ~0.5 for Nino3.4 Precipitation
 - ~0.1-0.2 for Z500 and CONUS T2m
 - <0.1 for CONUS Precipitation






Anomaly Correlations, Week 1



CFS_v2 raw CFS_v2 sec UFS_p1 raw UFS_p1 sec UFS_p2 raw UFS_p2 sec UFS_p3.1 raw UFS_p3.1 sec UFS_p4.0 raw UFS_p4.0 sec
UFS_p5.0 raw UFS_p5.0 sec






Anomaly Correlations, Week 2



CFS_v2 raw CFS_v2 sec UFS_p1 raw UFS_p1 sec UFS_p2 raw UFS_p2 sec UFS_p3.1 raw UFS_p3.1 sec UFS_p4.0 raw UFS_p4.0 sec
UFS_p5.0 raw UFS_p5.0 sec






Anomaly Correlations, Weeks 3&4



CFS_v2 raw CFS_v2 sec UFS_p1 raw UFS_p1 sec UFS_p2 raw UFS_p2 sec UFS_p3.1 raw UFS_p3.1 sec UFS_p4.0 raw UFS_p4.0 sec
UFS_p5.0 raw UFS_p5.0 sec






Anomaly Correlations, T2m, Week 1



P5


CPCTEMP UFSv5 SEC week 1

sec




P4

CPCTEMP UFSv4 SEC week 1




P3.1

CPCTEMP UFSv31 SEC week 1




CPCTEMP UFSv5 RAW week 1


raw



CPCTEMP UFSv4 RAW week 1



CPCTEMP UFSv31 RAW week 1



*CPC Global 0.5 degree Daily 2-m TMIN/TMAX from:
ftp://ftp.cpc.ncep.noaa.gov/precip/wd52ws/global_temp/
e.g., CPC_GLOBAL_T_V0.x_0.5deg.lnx.YYYY



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 10




Anomaly Correlations, T2m, Week 2



P5


CPCTEMP UFSv5 SEC week 2

sec




P4

CPCTEMP UFSv4 SEC week 2




P3.1

CPCTEMP UFSv31 SEC week 2




CPCTEMP UFSv5 RAW week 2


raw



CPCTEMP UFSv4 RAW week 2



CPCTEMP UFSv31 RAW week 2



*CPC Global 0.5 degree Daily 2-m T
<ftp://ftp.cpc.ncep.noaa.gov/precip/wd5>
e.g., CPC_GLOBAL_T_V0.x_0.5deg.inx.YYYY



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 11




Anomaly Correlations, T2m, Weeks 3&4



P5


CPCTEMP UFSv5 SEC week 34

sec




P4

CPCTEMP UFSv4 SEC week 34




P3.1

CPCTEMP UFSv31 SEC week 34




CPCTEMP UFSv5 RAW week 34


raw



CPCTEMP UFSv4 RAW week 34



CPCTEMP UFSv31 RAW week 34



*CPC Global 0.5 degree Daily 2-m T
<ftp://ftp.cpc.ncep.noaa.gov/precip/wd5>
e.g., CPC_GLOBAL_T_V0.x_0.5deg.inx.yyyr



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 12




Anomaly Correlations, Precipitation, Week 1



P5


CPCRAIN UFSv5 SEC week 1



sec


P4

CPCRAIN UFSv4 SEC week 1




P3.1

CPCRAIN UFSv31 SEC week 1




CPCRAIN UFSv5 RAW week 1





raw

CPCRAIN UFSv4 RAW week 1



CPCRAIN UFSv31 RAW week 1



*CPC Global 0.5 degree Unified Rain Gauge data from:
ftp://ftp.cpc.ncep.noaa.gov/precip/CPC_UNI_PRCP/GAUGE_GLB/
e.g., PRCP_CU_GAUGE_V1.0GLB_0.5deg.lnx.YYYYMMDDRT



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 13




Anomaly Correlations, Precipitation, Week 2




P5

CPCRAIN UFSv5 SEC week 2




P4

CPCRAIN UFSv4 SEC week 2




P3.1


CPCRAIN UFSv31 SEC week 2




CPCRAIN UFSv5 RAW week 2



CPCRAIN UFSv4 RAW week 2



CPCRAIN UFSv31 RAW week 2



*CPC Global 0.5 degree Unified Rain Gauge data from:
ftp://ftp.cpc.ncep.noaa.gov/precip/CPC_UNI_PRCP/GAUGE_GLB/
e.g., PRCP_CU_GAUGE_V1.0GLB_0.50deg.lnx.YYYYMMDDRT



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 14




Anomaly Correlations, Precipitation, Weeks 3&4



P5

CPCRAIN UFSv5 SEC week 34

sec



P4

CPCRAIN UFSv4 SEC week 34




P3.1

CPCRAIN UFSv31 SEC week 34



CPCRAIN UFSv5 RAW week 34

raw



CPCRAIN UFSv4 RAW week 34



CPCRAIN UFSv31 RAW week 34



*CPC Global 0.5 degree Unified Rain Gauge data from:
ftp://ftp.cpc.ncep.noaa.gov/precip/CPC_UNI_PRCP/GAUGE_GLB/
e.g., PRCP_CU_GAUGE_V1.0GLB_0.50deg.lnx.YYYYMMDDRT



NATIONAL WEATHER SERVICE

Building a Weather-Ready Nation // 15



MJO





Notes on the methodology used here



The methods to calculate the MJO index RMM1 & RMM2, and bivariate correlation skill are based on Wheeler and Hendon, 2004; Lin et al., 2008



The observations are the daily OLR from the CPC archives and U200 & U850 from CDAS2 in 01jan2010-30Sep2018



All-seasons MJO's two leading modes (RMM1 and RMM2) of the combined timeseries of OLR, U850 and U200 equatorial anomalies are shown. RMM1 series has the largest amplitude in the Maritime Continent and (negative) in the western hemisphere and Africa. RMM2 has largest amplitude in the Western Pacific and (negative) in the Indian Ocean.





Summary of findings




- **AC skill:**
 - To 0.6: ~16-18 days for combined RMM1+RMM2. Longer by ~3 days for RMM2 than for RMM1
 - To 0.5: ~21-24 days for combined RMM1+RMM2. Similar between RMM1 and RMM2
- **Amplitude error:** MJO is stronger than observations
- **Phase error:** MJO propagation is slower than observations






Anomaly Correlations, MJO: RMM1, RMM2, combined





MJO RMSE Skill




- $$RMSE(RMM1\&2) = \sqrt{\frac{1}{N} \sum_1^N [(RMM1(f) - RMM1(o))^2 + (RMM2(f) - RMM2(o))^2]}$$

N=1,168





MJO Amplitude and Phase Error



- $AMP = \sqrt{RMM1^2 + RMM2^2}$
- $E(AMP) = \frac{1}{N} \sum_1^N [AMP(f) - AMP(o)]$, $N=1,168$
- >0 means stronger; <0 means weaker than obs.

- MJO propagates too slowly

- $E(Phase) = \frac{1}{N} \sum_1^N \tan^{-1} \left(\frac{RMM1(o) * RMM2(f) - RMM2(o) * RMM1(f)}{RMM1(o) * RMM2(o) + RMM1(f) * RMM2(f)} \right)$ Kim et al (2018)
- >0 means faster; <0 means slower than obs.



Annual mean biases

Days 1-10, 11-20, 21-30












Summary of findings

- **SST:** Generally warm bias, particularly pronounced along ITCZ, Maritime continent, and western continental coasts. Cold bias in cold tongue region and near Gulfstream and Pacific ocean ~30N. Both warm and cold bias increase with lead time
- **Land Precipitation:** Generally wet bias, particularly strong in equatorial Africa and South America and southeast Asia. Dry bias in Indian subcontinent, northwestern Australia, and northern sub Saharan Africa
- **Land T2m:** Generally warm bias, with the exception of Central and South America and southeast Asia. For CONUS the warm bias affects the western US. Warm bias increases with lead time.
- **Z500:** Area-averaged bias is small and positive, somewhat increasing with lead time



SST bias



OSTIA SST used for verification














Precipitation bias (land)



CFSR used for verification












NATIONAL WEATHER SERVICE





T2m bias (land)












CFSR used for
verification





Z500 bias



CFSR used for
verification





Ocean flux biases

January/July



Summary of findings


- **SST:** Warm bias along ITCZ, Maritime continent, and western continental coasts. Tendency for warm bias at higher latitudes in winter hemisphere. Cold bias in cold tongue region.
- **Net Shortwave Radiation:** Similar pattern and sign to SST bias
- **Latent Heat Flux:** Generally positive bias, more pronounced along ITCZ, and around 40° latitude of the winter hemisphere. Particularly large in the North Atlantic in winter
- **Net Longwave Radiation:** Generally negative bias, especially pronounced in the western coasts of continents and high latitudes where SST has warm bias. No strong bias along ITCZ.
- **Sensible Heat Flux:** Positive bias in high latitudes of winter hemisphere: Kuroshio and Gulfstream in NH winter and in high southern latitudes in SH winter. No strong bias in the tropics
- **Wind Stress:** Positive bias in high latitudes of winter hemisphere and Arabian Sea in summer




P5 Jan/Jul (SST, netSW, LH)




SST: UFS P5-OSTIA JAN




net SW: UFS P5-OBS Jan




LH: UFS P5-ERA5 Jan







SST: UFS P5-OSTIA Jul



net SW: UFS P5-OBS Jul



LH: UFS P5-ERA5 Jul



P5 Jan/Jul (netLW, SH, Tau)

